

A Complete Classification of Complexity in Allen's Algebra in the Presence of a Non-Trivial Basic Relation

Andrei Krokhin* and Peter Jeavons*

Oxford University Computing Laboratory
Wolfson Building, Parks Road, OX1 3QD Oxford, UK
{andrei.krokhin, peter.jeavons}@comlab.ox.ac.uk

Peter Jonsson†

Department of Computer and Information Science
Linköping University, S-581 83 Linköping, Sweden
email: peter.jonsson@ida.liu.se

Abstract

We study fragments of Allen's algebra that contain a basic relation distinct from the equality relation. We prove that such a fragment is either NP-complete or else contained in some already known tractable subalgebra. We obtain this result by giving a new uniform description of known maximal tractable subalgebras and then systematically using an algebraic technique for description of maximal subalgebras with a given property. This approach avoids the need for extensive computer-assisted search.

1 Introduction

Allen's interval algebra [Allen, 1983] is one of the best established formalisms for temporal reasoning. Such reasoning is an important task in many areas of computer science and artificial intelligence (see, e.g., [Golumbic and Shamir, 1993; Nökel, 1991]). Allen's algebra allows us to specify qualitative information about time intervals. This algebra has also become the kernel of some other formalisms [Meiri, 1996; Angelsmark and Jonsson, 2000]. The interval algebra and some of its extensions are closely related with a number of interval-based temporal logics used for real-time system specification [Bellini *et al.*, 2000].

The basic satisfiability problem in Allen's algebra is NP-complete [Vilain *et al.*, 1989], so it is unlikely that efficient algorithms exist for reasoning in the full algebra. The computational difficulty has motivated the search for effective heuristics, e.g., [Ladkin and Reinefeld, 1992], and the study of the complexity of fragments of the algebra, e.g., [Golumbic and Shamir, 1993; Nebel and Bürckert, 1995].

*This research is partially supported by the UK EPSRC grants GR/R22704 and GR/M12926.

†This research is partially supported by the Swedish Research Council for Engineering Sciences (TFR) under grants 97-301 and 2000-361.

In this paper we follow the second approach, and we also assume throughout that $P \neq NP$. Research in this direction has focused on identifying *maximal* tractable fragments, i.e., fragments which cannot be extended without losing tractability. So far, eighteen maximal tractable fragments of the algebra have been identified [Drakengren and Jonsson, 1997a; 1997b; Nebel and Bürckert, 1995]. It is still unknown whether any others exist.

A complete classification of complexity within a certain large part of Allen's algebra was obtained in [Drakengren and Jonsson, 1998]. This result (as well as most similar results) was achieved by computer-assisted exhaustive search. However, it was noted in [Drakengren and Jonsson, 1998] that, for further progress, theoretical studies of the structure of Allen's algebra are required. There are some theoretical investigations of the structure of Allen's algebra, (see, e.g., [Hirsch, 1996; Ladkin and Maddux, 1994]). However they generally allow more operations on relations than originally used in [Allen, 1983], which makes them inappropriate for classifying complexity within the interval algebra. In this paper we systematically use an algebraic method that is similar to the approach taken in [Ligozat, 1998].

The first novel element in our approach is a new uniform description for all of the maximal tractable subalgebras of Allen's algebra which have already been identified (Table 2). Then we exhibit several new small NP-complete fragments of the algebra, including the first examples of NP-complete singletons (Proposition 2). Finally, we fully exploit the algebraic properties of Allen's algebra by importing a technique from general algebra. This technique has been used in many other contexts to obtain a description of maximal subalgebras of a given algebra with a given property (e.g., [Szendrei, 1995]). Here, for the first time, we systematically apply this technique to Allen's algebra to obtain a complete classification of the complexity of those fragments of the algebra containing a non-trivial basic relation (Theorem 1). Our method also provides (as a by-product) a new elementary proof of maximality of ten tractable subalgebras. In previous papers this proof of maximality has required extensive computer-aided case anal-

Basic relation		Example	Endpoints
x precedes y	p	xxx	$x^+ < y^-$
y preceded by x	p^{-1}	yyy	
x meets y	m	xxxx	$x^+ = y^-$
y met by x	m^{-1}	yyyy	
x overlaps y	o	xxxxx	$x^- < y^- < x^+$,
y overl. by x	o^{-1}	yyyyy	$x^+ < y^+$
x during y	d	xxx	$x^- > y^-$,
y includes x	d^{-1}	yyyyyyy	$x^+ < y^+$
x starts y	s	xxx	$x^- = y^-$,
y started by x	s^{-1}	yyyyyyy	$x^+ < y^+$
x finishes y	f	xxx	$x^+ = y^+$,
y finished by x	f^{-1}	yyyyyyy	$x^- > y^-$
x equals y	$\equiv$	xxxx yyyy	$x^- = y^-$, $x^+ = y^+$

Table 1: The thirteen basic relations. The endpoint relations $x^- < x^+$ and $y^- < y^+$ that are valid for all relations have been omitted.

ysis [Drakengren and Jonsson, 1997a; 1997b].

The paper is organized as follows: in Section 2 we give the basic definitions of Allen’s algebra and present the known maximal tractable subalgebras in the new form. Section 3 contains the new classification result, and Section 4 contains a brief discussion of future extensions.

2 Allen’s Algebra

Allen’s interval algebra [Allen, 1983] is based on the notion of *relations between intervals*. An interval x is represented as a tuple (x^-, x^+) of real numbers with $x^- < x^+$, denoting the left and right endpoints of the interval, respectively. The possible relations between intervals are the 2^{13} possible unions of 13 *basic interval relations*, which are shown in Table 1. Relations between intervals will also be written as collections of basic relations so, for instance, the union of the basic relations p, m and f^{-1} is written $(p \ m \ f^{-1})$. Allen’s algebra $\mathcal{A}$ consists of the 2^{13} possible relations between intervals together with the operations *converse* $\cdot^{-1}$, *intersection* $\cap$ and *composition* $\circ$ which are defined as follows:

$$\forall x, y : x r^{-1} y \Leftrightarrow y r x$$

$$\forall x, y : x(r \cap s)y \Leftrightarrow x r y \ \& \ x s y$$

$$\forall x, y : x(r \circ s)y \Leftrightarrow \exists z : (x r z \ \& \ z s y)$$

It follows that the converse of $r = (b_1 \dots b_n)$ is equal to $(b_1^{-1} \dots b_n^{-1})$. The intersection of two relations can be expressed as the usual set-theoretic intersection. Using the definition of composition, it can be derived that

$$(b_1 \dots b_n) \circ (b'_1 \dots b'_m) = \bigcup \{b_i \circ b'_j \mid i \leq n, j \leq m\}.$$

All algebraic calculations given in Section 3 can be done by hand; however, a simple table showing the results of the composition operation on all pairs of basic relations (which can be found, e.g., in [Ladkin and Maddux, 1994]) makes such calculations considerably easier.

The problem of *satisfiability* ($\mathcal{A}$ -SAT) of a set of interval variables with relations between them is that of deciding whether there exists an assignment of intervals on the real line for the interval variables, such that all of the relations between the intervals are satisfied. This is defined as follows.

Definition 1 Let $X \subseteq \mathcal{A}$ be a set of interval relations. An instance I of $\mathcal{A}$ -SAT(X) is a set, V , of variables and a set of constraints of the form $x r y$ where $x, y \in V$ and $r \in X$. The question is whether I is satisfiable, i.e., whether there exists a function, f , from V to the set of all intervals such that $f(x) r f(y)$ holds for every constraint $x r y$ in I .

If there exists a polynomial-time algorithm solving all instances of $\mathcal{A}$ -SAT(X) then we say that X is tractable. On the other hand, if $\mathcal{A}$ -SAT(X) is NP-complete then we say that X is NP-complete. Since the problem $\mathcal{A}$ -SAT($\mathcal{A}$) is NP-complete [Vilain *et al.*, 1989], there arises the question of description of tractable subsets of Allen’s algebra.

Subsets of $\mathcal{A}$ that are closed under the operations of intersection, converse and composition are said to be *subalgebras*. For a given subset X of $\mathcal{A}$, the smallest subalgebra containing X is called the subalgebra *generated* by X and is denoted by $\langle X \rangle$. It is easy to see that $\langle X \rangle$ is obtained from X by adding all relations that can be obtained from the relations in X by using the three operations of $\mathcal{A}$.

It is known [Nebel and Bürckert, 1995], and easy to prove, that, for every $X \subseteq \mathcal{A}$, the problem $\mathcal{A}$ -SAT($\langle X \rangle$) is polynomially equivalent to $\mathcal{A}$ -SAT(X). Therefore, to classify the complexity of all subsets of $\mathcal{A}$ it is only necessary to consider *subalgebras* of $\mathcal{A}$. Obviously, adding relations to a subalgebra can only increase the complexity of the corresponding satisfiability problem. Thus, since $\mathcal{A}$ is finite, the problem of describing tractability in $\mathcal{A}$ can be reduced to the problem of describing the *maximal* tractable subalgebras in $\mathcal{A}$, i.e., subalgebras that cannot be extended without losing tractability.

The known maximal tractable subalgebras [Drakengren and Jonsson, 1997a; 1997b; Nebel and Bürckert, 1995] are presented in Table 2. In our proofs as well as in Table 2 we use the symbol $\pm$, which should be interpreted as follows. A condition involving $\pm$ means the conjunction of two conditions: one corresponding to $+$ and one corresponding to $-$. For example, condition $(o)^{\pm} \subseteq r \Leftrightarrow (d)^{\pm} \subseteq r$ means that both $(o) \subseteq r \Leftrightarrow (d) \subseteq r$ and $(o^{-1}) \subseteq r \Leftrightarrow (d^{-1}) \subseteq r$ hold. The main advantage of using the $\pm$ symbol is conciseness: in any subalgebra of $\mathcal{A}$, the $+$ and the $-$ conditions are satisfied (or not satisfied) simultaneously, and, therefore only one of them needs to be verified.

In previous papers, the subalgebras from Table 2 were defined in other ways. However, in all cases except for $\mathcal{H}$, it is very straightforward to verify that our definitions are equivalent to the original ones. The subalgebra $\mathcal{H}$ was originally defined as the ‘ORD-Horn algebra’ [Nebel and Bürckert, 1995], but has also been characterized as the algebra of ‘pre-convex’ relations (see, e.g., [Ligozat, 1998]). Using the latter description it is not hard to show that our definition of $\mathcal{H}$ is equivalent. Finally, we note that there is one more known maximal tractable subalgebra that is not included in Table 2. That subalgebra is defined as $\{\emptyset\} \cup \{r \in \mathcal{A} \mid (\equiv) \subseteq r\}$ and the only basic relation it contains is $(\equiv)$.

$S_p = \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (p)^{\pm 1} \subseteq r\}$	$\mathcal{E}_p = \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (p)^{\pm 1} \subseteq r\}$
$S_d = \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (d^{-1})^{\pm 1} \subseteq r\}$	$\mathcal{E}_d = \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (d)^{\pm 1} \subseteq r\}$
$S_o = \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (o)^{\pm 1} \subseteq r\}$	$\mathcal{E}_o = \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (o)^{\pm 1} \subseteq r\}$
$A_1 = \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (s^{-1})^{\pm 1} \subseteq r\}$	$B_1 = \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (f^{-1})^{\pm 1} \subseteq r\}$
$A_2 = \{r \mid r \cap (\text{pmod}^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r\}$	$B_2 = \{r \mid r \cap (\text{pmods})^{\pm 1} \neq \emptyset \Rightarrow (f)^{\pm 1} \subseteq r\}$
$A_3 = \{r \mid r \cap (\text{pmod}f)^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r\}$	$B_3 = \{r \mid r \cap (\text{pmod}^{-1}s^{-1})^{\pm 1} \neq \emptyset \Rightarrow (f^{-1})^{\pm 1} \subseteq r\}$
$A_4 = \{r \mid r \cap (\text{pmod}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r\}$	$B_4 = \{r \mid r \cap (\text{pmod}^{-1}s)^{\pm 1} \neq \emptyset \Rightarrow (f^{-1})^{\pm 1} \subseteq r\}$
$\mathcal{E}^* = \left\{ r \mid \begin{array}{l} 1) r \cap (\text{pmod})^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r, \text{ and} \\ 2) r \cap (ff^{-1}) \neq \emptyset \Rightarrow (\equiv) \subseteq r \end{array} \right\}$	$\mathcal{S}^* = \left\{ r \mid \begin{array}{l} 1) r \cap (\text{pmod}^{-1})^{\pm 1} \neq \emptyset \Rightarrow (f^{-1})^{\pm 1} \subseteq r, \text{ and} \\ 2) r \cap (ss^{-1}) \neq \emptyset \Rightarrow (\equiv) \subseteq r \end{array} \right\}$
$\mathcal{H} = \left\{ r \mid \begin{array}{l} 1) r \cap (os)^{\pm 1} \neq \emptyset \ \& \ r \cap (o^{-1}f)^{\pm 1} \neq \emptyset \Rightarrow (d)^{\pm 1} \subseteq r, \text{ and} \\ 2) r \cap (ds)^{\pm 1} \neq \emptyset \ \& \ r \cap (d^{-1}f^{-1})^{\pm 1} \neq \emptyset \Rightarrow (o)^{\pm 1} \subseteq r, \text{ and} \\ 3) r \cap (pm)^{\pm 1} \neq \emptyset \ \& \ r \not\subseteq (pm)^{\pm 1} \Rightarrow (o)^{\pm 1} \subseteq r \end{array} \right\}$	

Table 2: Maximal tractable subalgebras of Allen's algebra.

3 Main Result

Throughout the paper, $\mathcal{S}$ denotes a subalgebra of $\mathcal{A}$. We denote by $\text{bas}(\mathcal{S})$ the set of basic relations in $\mathcal{S}$. Our result is an extension of the following proposition which was proved in [Drakengren and Jonsson, 1998].

Proposition 1

- 1) Let $\mathcal{S}$ be a subalgebra of $\mathcal{A}$ with $|\text{bas}(\mathcal{S})| > 3$. Then $\mathcal{S}$ is tractable if and only if it is contained in one of the following 7 algebras: $\mathcal{S}_p, \mathcal{S}_d, \mathcal{S}_o, \mathcal{E}_p, \mathcal{E}_d, \mathcal{E}_o$, and $\mathcal{H}$. Otherwise $\mathcal{S}$ is NP-complete.
- 2) Let $\mathcal{S}$ be a subalgebra of $\mathcal{A}$ such that $(m) \in \mathcal{S}$ or $(p) \in \mathcal{S}$. If $\mathcal{S}$ is tractable then $\mathcal{S} \subseteq \mathcal{S}_p$, or $\mathcal{S} \subseteq \mathcal{E}_p$, or $\mathcal{S} \subseteq \mathcal{H}$. Otherwise $\mathcal{S}$ is NP-complete.

We improve this result by showing that any so far unknown tractable subalgebra can contain only a *trivial* basic relation. Throughout the paper, by a *trivial* basic relation we mean the equality relation $\equiv$.

Theorem 1 Suppose $\mathcal{S}$ is a subalgebra of $\mathcal{A}$ containing a non-trivial basic relation. Then $\mathcal{S}$ is tractable if and only if it is contained in one of the 17 algebras listed in Table 2. Otherwise $\mathcal{S}$ is NP-complete.

We can assume, without loss of generality, that $\mathcal{S}$ contains the relation $(\equiv)$, since it is easy to show that $\mathcal{S}$ and $\mathcal{S} \cup \{(\equiv)\}$ have the same complexity (up to polynomial-time equivalence). This implies that the size of $\text{bas}(\mathcal{S})$ is odd since $\mathcal{S}$ is closed under converse. By combining Proposition 1(1) and the previous observation, we may consider only the case $|\text{bas}(\mathcal{S})| = 3$. By Proposition 1(2), it suffices to consider the cases where $\text{bas}(\mathcal{S})$ is one of the following sets: $\{\equiv, s, s^{-1}\}$, $\{\equiv, f, f^{-1}\}$, $\{\equiv, d, d^{-1}\}$, or $\{\equiv, o, o^{-1}\}$.

The rest of this section is organized as follows. In Subsection 3.1 we describe some NP-complete sets of relations which we use in our proofs later. In Subsection 3.2, we give a complete proof of Theorem 1 for the cases where $\text{bas}(\mathcal{S}) = \{\equiv, s, s^{-1}\}$ and $\text{bas}(\mathcal{S}) = \{\equiv, f, f^{-1}\}$. Finally, in Subsection 3.3 we give a sketch of a proof of Theorem 1 for the cases where $\text{bas}(\mathcal{S}) = \{\equiv, d, d^{-1}\}$ and $\text{bas}(\mathcal{S}) = \{\equiv, o, o^{-1}\}$.

Finally, we observe that when $\mathcal{S}$ is one of the algebras $\mathcal{E}^*$, $\mathcal{S}^*$, $\mathcal{A}_i$, or $\mathcal{B}_i$ ($1 \leq i \leq 4$), then $\text{bas}(\mathcal{S})$ is either $\{\equiv, s, s^{-1}\}$ or $\{\equiv, f, f^{-1}\}$. With the help of computer-aided exhaustive search, it was proved in [Drakengren and Jonsson, 1997a; 1997b] that these ten algebras are maximal tractable subalgebras of Allen's algebra. Since we do not use computer-aided results in Subsection 3.2, our results provide a new elementary proof of maximality for these algebras.

Throughout this section we shall assume that a non-trivial relation is also non-empty; this will cause no confusion.

3.1 NP-completeness results

This subsection contains the NP-completeness proofs that we need in the sequel. Our reductions are based on the NP-complete problem BETWEENNESS [Garey and Johnson, 1979]:

INSTANCE: A finite set A and a collection T of ordered triples (a, b, c) of distinct elements from A .

QUESTION: Is there a total ordering $<$ on A such that for each $(a, b, c) \in T$, we have either $a < b < c$ or $c < b < a$?

Let us fix relations $R, R_1, R_2 \in \mathcal{A}$. Then we define $\Gamma(a, b, c, x, y)$ to be the following problem instance over the variables $\{a, b, c, x, y\}$:

$$\{xR_1a, xR_1b, xR_2c, yR_2a, yR_1b, yR_1c\}.$$

Further, define $\Gamma_1 = \Gamma(a, b, c, x, y) \cup \{aRb, bRc\}$, $\Gamma_2 = \Gamma(a, b, c, x, y) \cup \{bRa, bRc, aR \cup R^{-1}c\}$, and $\Gamma_3 = \Gamma(a, b, c, x, y) \cup \{aRb, cRb, aR \cup R^{-1}c\}$.

Lemma 1 Let $R \in \{(p), (d), (o), (s)\}$, $R_1, R_2 \in \mathcal{A}$, and let $\Gamma_1, \Gamma_2, \Gamma_3$ be as above. If Γ_1 is satisfiable while Γ_2 and Γ_3 are not, then $\{R \cup R^{-1}, R_1, R_2\}$ is NP-complete.

Proof. Polynomial-time reduction from BETWEENNESS. Let (A, T) be an arbitrary instance of the problem and construct an instance I of $\mathcal{A}\text{-SAT}(\{R \cup R^{-1}, R_1, R_2\})$ as follows:

- (1) for each pair of distinct elements $a, b \in A$, add the constraint $aR \cup R^{-1}b$ to I ; and

- (2) for each triple $(a, b, c) \in T$, introduce two fresh variables x, y and add $\Gamma(a, b, c, x, y)$ to I .

It is a routine verification to show that I is satisfiable if and only if (A, T) has a solution. $\square$

The restriction $R \in \{(p), (d), (o), (s)\}$ is imposed to ensure that we get a natural total ordering in the reduction. Many other relations such as (dsf) have this property but there are also relations without this property, e.g., $(\equiv)$ and (mm^{-1}) .

Suppose $X \subseteq \mathcal{A}$ and I is an instance of $\mathcal{A}\text{-SAT}(X)$. Let variables x, y be involved in I . Further, let $r \in \mathcal{A}$ be the relation defined as follows. A basic relation r' is included in r if and only if the instance obtained from I by adding the constraint $xx'y$ is satisfiable. In this case, we say that r is *derived* from X . It can be easily checked that the problems $\mathcal{A}\text{-SAT}(X)$ and $\mathcal{A}\text{-SAT}(X \cup \{r\})$ are polynomially equivalent. Looking now at the definition of the three operations of the interval algebra, it is easy to see that, generally, one can derive more relations from a given $X \subseteq \mathcal{A}$ than one can generate using these operations. On the other hand, derivation is essentially harder to manage while the operations of Allen's algebra give us the advantage of employing algebraic techniques. Therefore, in this paper, we use derivation only once in the proof of the following statement.

Proposition 2 *The following subsets of $\mathcal{A}$ are NP-complete:*

- 1) $\{(d), (oo^{-1})\}$, $\{(d^{-1}), (pp^{-1})\}$, and $\{(o), (dd^{-1})\}$.
- 2) $\{r\}$ with $(ods^{-1}) \subseteq r \subseteq (pmod s^{-1}f^{-1})$.

Proof. For part (1), apply Lemma 1 with $R = (o)$, $R_1 = (d)$, $R_2 = (oo^{-1})$, or with $R = (p)$, $R_1 = (d^{-1})$, $R_2 = (pp^{-1})$, or with $R = (d)$, $R_1 = (o)$, $R_2 = (dd^{-1})$, respectively.

To prove part (2), let r_3 be the union of all basic relations except for $\equiv$ and s^{-1} , and consider the instance $\Gamma_4 = \{arb, brc, cra, drb, drc\}$ over the variables a, b, c, d . In the cases when $r = (ods^{-1})$ or $r = (pmod s^{-1}f^{-1})$, it can be shown that $\Gamma_4 \cup \{dr'a\}$ is satisfiable for every basic relation $r' \subseteq r_3$ but not satisfiable for any other choice of r' . Hence, we can derive r_3 from r , and we can also derive the relations $r_4 = r \cap r_3 = r - (s^{-1})$ and $r_5 = r_4 \circ r_4$. It is easy to check that

$$(pmod s) \subseteq r_5 \subseteq (pmod s f^{-1})$$

which implies that $r_5 \cap r^{-1} = (s)$. Furthermore, we have $(s) \circ (s^{-1}) = (\equiv ss^{-1})$, and $r \circ r$ is the disequality relation, so the relation (ss^{-1}) can be obtained from the relation r . If $R = (s)$, $R_1 = r^{-1}$ and $R_2 = r$, then these relations satisfy the conditions of Lemma 1 so $\{(ss^{-1}), r^{-1}, r\}$ is NP-complete. Since all of these relations can be derived from the single relation r , it follows that $\{r\}$ is NP-complete. $\square$

3.2 The s/f case

We will show that if $bas(\mathcal{S}) = \{\equiv, s, s^{-1}\}$, then either $\mathcal{S}$ is NP-complete or $\mathcal{S}$ is contained in one of the subalgebras $\mathcal{S}_p, \mathcal{S}_d, \mathcal{S}_o, \mathcal{E}^*, \mathcal{H}$, or in one of $\mathcal{A}_i$, $1 \leq i \leq 4$. By using the obvious symmetry between the relations (s) and (f) , it immediately follows that if $bas(\mathcal{S}) = \{\equiv, f, f^{-1}\}$, then either $\mathcal{S}$ is NP-complete or contained in one of $\mathcal{E}_p, \mathcal{E}_d, \mathcal{E}_o, \mathcal{S}^*, \mathcal{H}$,

or $\mathcal{B}_i$, $1 \leq i \leq 4$. For the remainder of this subsection, it is assumed that $bas(\mathcal{S}) = \{\equiv, s, s^{-1}\}$.

Given a relation r , we write r^* to denote the relation $r \cap r^{-1}$. Throughout the proofs we use the obvious fact that if $r_1 \subseteq r_2$ then, for any r , we have $r \circ r_1 \subseteq r \circ r_2$ and $r_1 \circ r \subseteq r_2 \circ r$. If b is a basic relation then r_b denotes the least relation $r \in \mathcal{S}$ such that $(b) \subseteq r$, i.e. the intersection of all $r \in \mathcal{S}$ with this property. It is easy to show that the relation r_b exists in $\mathcal{S}$ for every basic relation b , except for two very simple cases: when every $r \in \mathcal{S}$ satisfies exactly one of conditions $r \subseteq (\equiv ss^{-1})$ and $r \subseteq (\equiv ff^{-1})$. In both cases, we have $\mathcal{S} \subseteq \mathcal{H}$.

Lemma 2 *If $\mathcal{S}$ contains the relation (od) , then every relation in $\mathcal{S}$ satisfies condition 3) of $\mathcal{H}$.*

Proof. Arbitrarily choose $r \in \mathcal{S}$. Since $(od) \in \mathcal{S}$, it follows that $r_d = r_o = (od)$ and $(o)^{\pm 1} \subseteq r \Leftrightarrow (d)^{\pm 1} \subseteq r$. Furthermore, $r_1 = (s) \circ (od) = (pmod) \in \mathcal{S}$ so $r_p \subseteq r_1$ and $r_m \subseteq r_1$. Since $(pm) \circ (pm) = (p)$, we have $r_p \not\subseteq (pm)$ and $r_m \not\subseteq (pm)$, which implies that $r_p \cap (od) \neq \emptyset$ and $r_m \cap (od) \neq \emptyset$. Now it follows that $(od) \subseteq r_p$ and $(od) \subseteq r_m$. Thus, if $r \cap (pm) \neq \emptyset$, then $(od) \subseteq r$ and r satisfies condition 3) of $\mathcal{H}$. $\square$

Lemma 3 *If $\mathcal{S}$ contains a non-trivial relation r' with $r' \subseteq (\equiv pp^{-1}mm^{-1}ff^{-1})$ then $\mathcal{S}$ is included in one of $\mathcal{H}, \mathcal{E}^*$, or $\mathcal{S}_p$, or else $\mathcal{S}$ is NP-complete.*

Proof. Case 1. $r' = (ff^{-1})$

Since $(ff^{-1}) \circ (s) = (od) \in \mathcal{S}$, it follows that any $r \in \mathcal{S}$ satisfies condition 3) of $\mathcal{H}$, by Lemma 2. Now it follows that, for any $r \in \mathcal{S}$, we have $(o)^{\pm 1} \subseteq r \Leftrightarrow (d)^{\pm 1} \subseteq r$ and also $(f) \subseteq r \Leftrightarrow (f^{-1}) \subseteq r$.

Suppose that $\mathcal{S} \not\subseteq \mathcal{H}$, i.e. some $r \in \mathcal{S}$ fails to satisfy condition 1) or condition 2) of $\mathcal{H}$. Then, using the facts from the previous paragraph, it can be shown that we can choose r so that $(s^{-1}ff^{-1}) \subseteq r \subseteq (\equiv pmod ss^{-1}ff^{-1})$ or $(ods^{-1}) \subseteq r \subseteq (\equiv pmod ss^{-1}ff^{-1})$. In both cases, composing the relations with (s) from the left we get

$$(\equiv pmod ss^{-1}) \subseteq (s) \circ r \subseteq (\equiv pmod ss^{-1}).$$

Therefore $(\equiv pmod ss^{-1}) \in \mathcal{S}$, and $(pmod s^{-1})$ can be obtained as

$$(\equiv pmod ss^{-1}) \cap ((\equiv pmod ss^{-1})^{-1} \circ (od)^{-1}).$$

Then $\mathcal{S}$ is NP-complete, by Proposition 2(2).

Case 2. $r' \subseteq (\equiv ff^{-1})$.

Composing r' and its inverse we get $(\equiv ff^{-1})$, so we may assume that $r' = (\equiv ff^{-1})$. If some relation $r_2 \in \mathcal{S}$ fails to satisfy condition 2) of $\mathcal{E}^*$, then $r_2 \cap r'$ is either one of (f) and (f^{-1}) , which is impossible under our assumptions, or (ff^{-1}) going back to Case 1. Suppose now that each $r \in \mathcal{S}$ satisfies condition 2) of $\mathcal{E}^*$.

We have $r' \circ (s) = (ods) \in \mathcal{S}$. If (od) belongs to $\mathcal{S}$ then $r' \cap ((s^{-1}) \circ (od)) = (ff^{-1}) \in \mathcal{S}$, which implies $\mathcal{S} \subseteq \mathcal{H}$. Otherwise we have $r \cap (od)^{\pm 1} \neq \emptyset \Rightarrow (s)^{\pm 1} \subseteq r$ for every $r \in \mathcal{S}$. Assume a relation $r_3 \in \mathcal{S}$ does not satisfy condition 1) of $\mathcal{E}^*$, that is $r_3 \cap (pmod) \neq \emptyset$ and $(s) \not\subseteq r_3$. Since $(pmod s) = (s) \circ r' \in \mathcal{S}$, we have $r_4 = r_3 \cap (pmod s) \in \mathcal{S}$

and $r_4 \subseteq (\text{pm})$. This implies $r_4 \circ r_4 = (\text{p})$, a contradiction. Therefore the relations in $\mathcal{S}$ must satisfy both conditions of $\mathcal{E}^*$, that is $\mathcal{S} \subseteq \mathcal{E}^*$.

Case 3. $r' \subseteq (\equiv \text{mm}^{-1}\text{ff}^{-1})$ and $r' \cap (\text{mm}^{-1}) \neq \emptyset$.

Without loss of generality we may assume that $(\text{m}) \subseteq r'$. Then the relation $(r' \cap (r' \circ (\text{s}^{-1})))$ is equal either to (mm^{-1}) or to (m) . In the former case it follows that the relation $(\text{m}) = (\text{mm}^{-1}) \cap ((\text{mm}^{-1}) \circ (\text{s}^{-1}))$ belongs to $\mathcal{S}$ as well. A contradiction.

Case 4. $r' \cap (\text{pp}^{-1}) \neq \emptyset$.

We may assume, without loss of generality, that $(\text{p}) \subseteq r'$. Then $(\text{p}) \subseteq (\text{s}) \circ r' \subseteq (\text{pp}^{-1}\text{mm}^{-1}\text{ods})$. Furthermore, $(\text{p}) \subseteq r_5 = r' \cap ((\text{s}) \circ r') \subseteq (\text{pp}^{-1}\text{mm}^{-1})$, and we have $(\text{p}) \subseteq r_6 = r_5 \circ (\text{s}^{-1}) \subseteq (\text{pp}^{-1}\text{m})$. If $(\text{p}^{-1}) \not\subseteq r_6$ then $r_6 \circ r_6 = (\text{p}) \in \mathcal{S}$, a contradiction. Otherwise we have $(\text{pp}^{-1}) = r_6^* \in \mathcal{S}$. Then $(\text{p}) \subseteq r \Leftrightarrow (\text{p}^{-1}) \subseteq r$ holds for every $r \in \mathcal{S}$.

We have $(\text{s}^{-1}) \circ (\text{pp}^{-1}) = (\text{pp}^{-1}\text{mod}^{-1}\text{f}^{-1}) \in \mathcal{S}$. For every non-empty $r_7 \subseteq (\text{modf}^{-1})$, we have $((\text{s}) \circ r_7) \cap (\text{pp}^{-1}) = (\text{p})$. Therefore no such r_7 belongs to $\mathcal{S}$. We conclude that, for any $r \in \mathcal{S}$, if $r \cap (\text{mod}^{-1}\text{f}^{-1}) \neq \emptyset$ then $(\text{p}) \subseteq r$, which means that $\mathcal{S} \subseteq \mathcal{S}_\text{p}$. $\square$

We assume further that, for every non-trivial $r \in \mathcal{S}$, we have $r \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) \neq \emptyset$. In the next four lemmas we consider the case when $\mathcal{S}$ contains a non-trivial relation r' with $r' \cap (\text{ss}^{-1}) = \emptyset$. In view of the assumption just made we may consider that r' is either r_d or r_o . It is not hard to verify that these four lemmas indeed cover all possible cases for r_d and r_o .

Lemma 4 Suppose $r_\text{d} \cap (\text{ss}^{-1}) = \emptyset$. If $r_\text{d} \cap (\text{oo}^{-1}\text{dd}^{-1}) = (\text{d})$ or $(\text{dd}^{-1}) \subseteq r_\text{d}$ then $\mathcal{S} \subseteq \mathcal{S}_\text{d}$.

Proof. **Case 1.** $r_\text{d} \cap (\text{oo}^{-1}\text{dd}^{-1}) = (\text{d})$.

By assumption, the relation r_d satisfies condition (d) $\subseteq r_\text{d} \subseteq (\equiv \text{pp}^{-1}\text{mm}^{-1}\text{dff}^{-1})$. Let r_1 be calculated as $r_\text{d} \cap ((\text{s}) \circ r_\text{d})$. Then we have $(\text{d}) \subseteq r_1 \subseteq (\text{pp}^{-1}\text{mm}^{-1}\text{d})$. By minimality of r_d we get $(\text{d}) \subseteq r_\text{d} \subseteq (\text{pp}^{-1}\text{mm}^{-1}\text{d})$. Calculating r_1 again, we get $(\text{d}) \subseteq r_\text{d} \subseteq (\text{pp}^{-1}\text{m}^{-1}\text{d})$. Since $(\text{pp}^{-1}) \not\subseteq \mathcal{S}$, and since we have $(\text{m}^{-1}\text{d}) \subseteq r_\text{d} \circ (\text{s})$ only if $(\text{p}^{-1}) \subseteq r_\text{d}$, we may assume that r_d is either (pd) , or (p^{-1}d) , or $(\text{p}^{-1}\text{m}^{-1}\text{d})$. The first case is impossible because of $(\text{pd}) \cap ((\text{p}^{-1}\text{d}^{-1}) \circ (\text{s})) = (\text{d})$. Let $(\text{p}^{-1}\text{d}) \subseteq r_\text{d} \subseteq (\text{p}^{-1}\text{m}^{-1}\text{d})$. Then $(\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{df}) = r_\text{d} \circ (\text{s}) \in \mathcal{S}$. Suppose $\mathcal{S}$ contains a non-empty subrelation r_2 of $(\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{f})$. Then $r_3 = r_\text{d} \cap (r_2 \circ (\text{s}^{-1}))$ is a non-empty subrelation of $(\text{p}^{-1}\text{m}^{-1})$ implying that $(\text{p}^{-1}) = r_3 \circ r_3 \in \mathcal{S}$, a contradiction. Therefore, for every $r \in \mathcal{S}$, we have $r \cap (\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{f}) \neq \emptyset \Rightarrow (\text{d}) \subseteq r$, which means that $\mathcal{S} \subseteq \mathcal{S}_\text{d}$.

Case 2. $(\text{dd}^{-1}) \subseteq r_\text{d} \cap (\text{oo}^{-1}\text{dd}^{-1})$.

Suppose a relation $r_4 \in \mathcal{S}$ satisfies both $r_4 \not\subseteq (\equiv \text{ss}^{-1})$ and $(\text{dd}^{-1}) \not\subseteq r_4$. Then it is not hard to verify that either $((\text{s}^{-1}) \circ r_4) \cap r_\text{d}$ or $(r_4 \circ (\text{s})) \cap r_\text{d}$ contains exactly one of (d) and (d^{-1}) which contradicts the minimality of r_d . Thus, for every $r \in \mathcal{S}$ such that $r \not\subseteq (\equiv \text{ss}^{-1})$, we have $(\text{dd}^{-1}) \subseteq r$. This implies that $\mathcal{S} \subseteq \mathcal{S}_\text{d}$. $\square$

Lemma 5 Suppose $r_\text{o} \cap (\text{ss}^{-1}) = \emptyset$. If $r_\text{o} \cap (\text{oo}^{-1}\text{dd}^{-1}) = (\text{o})$ or $(\text{oo}^{-1}) \subseteq r_\text{o}$ then $\mathcal{S} \subseteq \mathcal{S}_\text{o}$.

Proof. Similar to the previous lemma. $\square$

Lemma 6 If $r_\text{d} = r_\text{o}^{-1}$ and $r_\text{d} \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) = (\text{o}^{-1}\text{d})$ then $\mathcal{S} \subseteq \mathcal{S}_\text{d}$.

Proof. As in the proof of Lemma 4, we can obtain $(\text{o}^{-1}\text{d}) \subseteq r_\text{d} \subseteq (\text{pp}^{-1}\text{m}^{-1}\text{o}^{-1}\text{df})$. If $(\text{po}^{-1}\text{d}) \subseteq r_\text{d}$ then $(r_\text{d} \circ (\text{s}^{-1}))^* = (\text{pp}^{-1}) \in \mathcal{S}$, a contradiction. Therefore we have $(\text{o}^{-1}\text{d}) \subseteq r_\text{d} \subseteq (\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{df})$, and $(\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{df}) = r_\text{d} \circ (\text{s}^{-1}) \in \mathcal{S}$. If some non-empty subrelation r_1 of $(\text{p}^{-1}\text{m}^{-1}\text{f})$ belongs to $\mathcal{S}$ then $r_2 = r_1 \cap (r_1 \circ (\text{s}^{-1}))$ is a non-empty subrelation of $(\text{p}^{-1}\text{m}^{-1})$. Then $r_2 \circ r_2 = (\text{p}^{-1})$ which contradicts our assumptions. Therefore, for every $r \in \mathcal{S}$, we have $r \cap (\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{f}) \neq \emptyset \Rightarrow (\text{d}) \subseteq r$, which means that $\mathcal{S} \subseteq \mathcal{S}_\text{d}$. $\square$

Lemma 7 If $r_\text{d} = r_\text{o}$ and $r_\text{d} \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ss}^{-1}) = (\text{od})$ then $\mathcal{S} \subseteq \mathcal{H}$ or $\mathcal{S}$ is NP-complete.

Proof. As in the previous lemmas, it can be shown that $(\text{od}) \subseteq r_\text{d} \subseteq (\text{pp}^{-1}\text{mm}^{-1}\text{od})$. It follows that $r_\text{d}^* = \emptyset$ and $(\text{pmod}) = (\text{s}) \circ r_\text{d} \in \mathcal{S}$. Furthermore, $(\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1}) = ((\text{s}^{-1}) \circ (\text{pmod}))^* \in \mathcal{S}$ and $(\text{od}) = (\text{pmod}) \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1}) \in \mathcal{S}$. By Lemma 2, we know that every $r \in \mathcal{S}$ satisfies condition 3) of $\mathcal{H}$. Suppose some $r_1 \in \mathcal{S}$ does not satisfy condition 1) of $\mathcal{H}$. Then r_1 can be chosen so that $(\text{s}^{-1}\text{f}^{-1}) \subseteq r_1 \subseteq (\equiv \text{pmodss}^{-1}\text{ff}^{-1})$ or $(\text{os}^{-1}) \subseteq r_1 \subseteq (\equiv \text{pmodss}^{-1}\text{ff}^{-1})$. If $(\text{f}) \subseteq r_1$ then $(\text{ff}^{-1}) = (r_1 \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1}))^* \in \mathcal{S}$ which contradicts our assumptions. Further, if $(\text{od}) \not\subseteq r_1$ then $r_1 \cap (\text{oo}^{-1}\text{dd}^{-1}\text{ff}^{-1}) = (\text{f}^{-1}) \in \mathcal{S}$, a contradiction. Therefore we may assume that $(\text{ods}^{-1}\text{f}^{-1}) \subseteq r_1 \subseteq (\equiv \text{pmodss}^{-1}\text{ff}^{-1})$. Now it can be checked that $r_2 = r_1 \cap (r_1^{-1} \circ (\text{o}^{-1}\text{d}^{-1}))$ satisfies $(\text{ods}^{-1}\text{f}^{-1}) \subseteq r_2 \subseteq (\text{pmodss}^{-1}\text{ff}^{-1})$. Then $\{r_2\}$ is NP-complete by Proposition 2(2). One can proceed similarly if condition 2) of $\mathcal{H}$ fails in $\mathcal{S}$. $\square$

It remains to consider the case when each non-trivial $r \in \mathcal{S}$ satisfies $r \cap (\text{ss}^{-1}) \neq \emptyset$.

Lemma 8 If $r \cap (\text{ss}^{-1}) \neq \emptyset$ for any non-trivial $r \in \mathcal{S}$ then $\mathcal{S} \subseteq \mathcal{A}_i$ for some $1 \leq i \leq 4$.

Proof. **Case 1.** $r_\text{p} \cap (\text{ss}^{-1}) = (\text{s}^{-1})$.

We have $(\text{ps}^{-1}) \subseteq r_\text{p}$ and $(\text{s}) \not\subseteq r_\text{p}$. Let $r_1 = (\text{s}^{-1}) \circ r_\text{p} \in \mathcal{S}$. Then it is easy to check that $(\text{pmod}^{-1}\text{s}^{-1}\text{f}^{-1}) \subseteq r_1$, and that $r_1 \cap (\equiv \text{s}) = \emptyset$. It follows that $r_1 = (\text{pmod}^{-1}\text{s}^{-1}\text{f}^{-1})$, since otherwise r_1^* is non-empty and $r_1^* \cap (\text{ss}^{-1}) = \emptyset$. No non-empty subrelation of $(\text{pmod}^{-1}\text{f}^{-1})$ can belong to $\mathcal{S}$. Therefore, for any $r \in \mathcal{S}$, $r \cap (\text{pmod}^{-1}\text{f}^{-1}) \neq \emptyset$ implies $(\text{s}^{-1}) \subseteq r$. Hence $\mathcal{S} \subseteq \mathcal{A}_1$.

Case 2. $r_\text{d} \cap (\text{ss}^{-1}) = (\text{s}^{-1})$.

The proof is similar to Case 1; the only change is that $r_1 = r_\text{d} \circ (\text{s}^{-1})$, and we deduce that $r_1 = (\text{p}^{-1}\text{m}^{-1}\text{o}^{-1}\text{ds}^{-1}\text{f})$, and, hence, $\mathcal{S} \subseteq \mathcal{A}_2$.

Case 3. $r_\text{o} \cap (\text{ss}^{-1}) = (\text{s}^{-1})$.

In view of Cases 1 and 2 we may assume that $(\text{os}^{-1}) \subseteq r_\text{o} \subseteq (\equiv \text{p}^{-1}\text{mm}^{-1}\text{od}^{-1}\text{s}^{-1}\text{ff}^{-1})$. Let $r_1 = (\text{s}^{-1}) \circ r_\text{o}$. It is easy to check that

$$(\text{od}^{-1}\text{s}^{-1}\text{f}^{-1}) \subseteq r_1 \subseteq (\text{p}^{-1}\text{m}^{-1}\text{oo}^{-1}\text{d}^{-1}\text{s}^{-1}\text{f}^{-1}).$$

Since $r_1^* \neq (oo^{-1})$, we obtain $r_1 \subseteq (p^{-1}m^{-1}od^{-1}s^{-1}f^{-1})$. It can straightforwardly be verified that if $r_1 \neq (od^{-1}s^{-1}f^{-1})$, then $r_2 = (r_1 \circ r_1)^* \in \mathcal{S}$, but $r_2 \cap (pp^{-1}ss^{-1}) = (pp^{-1})$, which contradicts the assumption made. Therefore, $r_1 = (od^{-1}s^{-1}f^{-1})$, and $(pmod^{-1}s^{-1}f^{-1}) = r_1 \circ r_1 \in \mathcal{S}$. Hence, for any $r \in \mathcal{S}$, $r \cap (pmod^{-1}f^{-1}) \neq \emptyset$ implies $(s^{-1}) \subseteq r$, and we have $\mathcal{S} \subseteq \mathcal{A}_1$.

Case 4. $r_m \cap (ss^{-1}) = (s^{-1})$.

Similarly to Case 3, we infer that $\mathcal{S} \subseteq \mathcal{A}_1$.

Case 5. (s) is contained in each of r_p, r_d, r_o , and r_m .

We have $r_f \cap (ss^{-1}) \neq \emptyset$. Then it follows that if $(s) \subseteq r_f$ then $\mathcal{S} \subseteq \mathcal{A}_3$. Otherwise, $(s^{-1}) \subseteq r_f$ and we have $\mathcal{S} \subseteq \mathcal{A}_4$. $\square$

3.3 The d/o case

Suppose $bas(\mathcal{S}) = \{\equiv, d, d^{-1}\}$. We give a sketch of a proof that either $\mathcal{S}$ is contained in $\mathcal{S}_d$ or in $\mathcal{E}_d$ (and therefore is tractable), or else it contains (oo^{-1}) or (pp^{-1}) (and then it is NP-complete by Proposition 2(1)).

Let $sym(\mathcal{S}) = \{r^* | r \in \mathcal{S}\}$. First it can be shown that either $sym(\mathcal{S})$ is contained in $\mathcal{S}_d$ or in $\mathcal{E}_d$, or else $\mathcal{S}$ contains (oo^{-1}) or (pp^{-1}) . Then one can check, similarly to Lemma 3, that if there exists a non-trivial $r \in \mathcal{S}$ such that $r \cap (dd^{-1}) = \emptyset$ then r satisfies $r \subseteq (\equiv ss^{-1})$ or $r \subseteq (\equiv ff^{-1})$, and in this case, once again, $\mathcal{S}$ is contained in $\mathcal{S}_d$ or in $\mathcal{E}_d$, respectively, or else $\mathcal{S}$ contains (oo^{-1}) or (pp^{-1}) . Finally, we assume that each non-trivial $r \in \mathcal{S}$ satisfies $r \cap (dd^{-1}) \neq \emptyset$, and then the required result can be proved by analysing the minimal relations r_p, r_m, r_o, r_s , and r_f , as in the proof of Lemma 8.

The case $bas(\mathcal{S}) = \{\equiv, o, o^{-1}\}$ is very similar to the previous one. The only essential difference is that the relation (pp^{-1}) cannot belong to $\mathcal{S}$; otherwise we get $(o) \circ (o) = (pmo)$ and $(p) = (pp^{-1}) \cap (pmo) \in \mathcal{S}$, a contradiction.

4 Conclusion

We have proved that any so far unknown tractable fragment of Allen's algebra can contain no other basic relation than $(\equiv)$. We believe that, by extending the algebraic techniques used to establish our result, it will be possible to show that there are in fact no other forms of tractability in Allen's algebra. In other words, we conjecture that all maximal tractable subalgebras are already identified. However, the proof of this conjecture requires a more elaborate algebraic argument than the proofs in this paper, since in the general case we have no particular relation to start with. Algebraic techniques similar to the one used in this paper can also be applied to the study of the complexity in other temporal and spatial formalisms.

References

[Allen, 1983] James F. Allen. Maintaining knowledge about temporal intervals. *Communications of the ACM*, 26(11):832–843, 1983.

[Angelsmark and Jonsson, 2000] Ola Angelsmark and Peter Jonsson. Some observations on durations, scheduling and Allen's algebra. In *Proceedings of the 6th Conference on Constraint Programming (CP'00)*, volume 1894 of *Lecture Notes in Computer Science*, pages 484–488. Springer-Verlag, 2000.

[Bellini et al., 2000] P. Bellini, R. Mattolini, and P. Nesi. Temporal logics for real-time system specification. *ACM Computing Surveys*, 32(1):12–42, 2000.

[Drakengren and Jonsson, 1997a] Thomas Drakengren and Peter Jonsson. Eight maximal tractable subclasses of Allen's algebra with metric time. *Journal of Artificial Intelligence Research*, 7:25–45, 1997.

[Drakengren and Jonsson, 1997b] Thomas Drakengren and Peter Jonsson. Twenty-one large tractable subclasses of Allen's algebra. *Artificial Intelligence*, 93(1-2):297–319, 1997.

[Drakengren and Jonsson, 1998] Thomas Drakengren and Peter Jonsson. A complete classification of tractability in Allen's algebra relative to subsets of basic relations. *Artificial Intelligence*, 106(2):205–219, 1998.

[Garey and Johnson, 1979] Michael Garey and David Johnson. *Computers and Intractability: A Guide to the Theory of NP-Completeness*. Freeman, New York, 1979.

[Golumbic and Shamir, 1993] Martin C. Golumbic and Ron Shamir. Complexity and algorithms for reasoning about time: A graph-theoretic approach. *Journal of the ACM*, 40(5):1108–1133, 1993.

[Hirsch, 1996] Robin Hirsch. Relation algebras of intervals. *Artificial Intelligence*, 83(2):1–29, 1996.

[Ladkin and Maddux, 1994] Peter B. Ladkin and Roger D. Maddux. On binary constraint problems. *Journal of the ACM*, 41(3):435–469, 1994.

[Ladkin and Reinefeld, 1992] Peter B. Ladkin and Alexander Reinefeld. Effective solution of qualitative interval constraint problems. *Artificial Intelligence*, 57(1):105–124, 1992.

[Ligozat, 1998] Gérard Ligozat. “Corner” relations in Allen's algebra. *Constraints*, 3(2-3):165–177, 1998.

[Meiri, 1996] Itay Meiri. Combining qualitative and quantitative constraints in temporal reasoning. *Artificial Intelligence*, 87(1-2):343–385, 1996.

[Nebel and Bürckert, 1995] Bernhard Nebel and Hans-Jürgen Bürckert. Reasoning about temporal relations: A maximal tractable subclass of Allen's interval algebra. *Journal of the ACM*, 42(1):43–66, 1995.

[Nökel, 1991] Klaus Nökel. *Temporally Distributed Symptoms in Technical Diagnosis*, volume 517 of *Lecture Notes in Artificial Intelligence*. Springer-Verlag, 1991.

[Szendrei, 1995] Ágnes Szendrei. Maximal non-affine reducts of simple affine algebras. *Algebra Universalis*, 34(1):144–174, 1995.

[Vilain et al., 1989] Marc B. Vilain, Henry A. Kautz, and Peter G. van Beek. Constraint propagation algorithms for temporal reasoning: A revised report. In Daniel S. Weld and Johan de Kleer, editors, *Readings in Qualitative Reasoning about Physical Systems*, pages 373–381. Morgan Kaufmann, San Mateo, CA, 1989.